WU B 02-08, PITHA 02/18, hep-ph/0212138

Proton mass effects in wide-angle Compton scattering

M. Diehl, Th. Feldmann
Institut für Theoretische Physik E, RWTH Aachen, 52056 Aachen, Germany

H.W. Huang
Department of Physics, University of Colorado, Boulder, CO 80309-0390, USA

P. Kroll
Fachbereich Physik, Universität Wuppertal, 42097 Wuppertal, Germany
(Dated: December 2002)

We investigate proton mass effects in the handbag approach to wide-angle Compton scattering. We find that theoretical uncertainties due to the proton mass are significant for photon energies presently studied at Jefferson Lab. With the proposed energy upgrade such uncertainties will be clearly reduced.

PACS numbers: 13.60.Fz

arXiv:hep-ph/0212138v1 10 Dec 2002

In Refs. [1–3] we have investigated the handbag approach to wide-angle Compton scattering off protons, $\gamma p \to \gamma p$. Analogous results have been obtained in Ref. [4]. In the handbag approach the Compton amplitude is given by a hard scattering $\gamma q \to \gamma q$ at parton level multiplied by soft Compton form factors describing the emission and reabsorption of the quark by the proton. The kinematical requirement for the applicability of this approach is that the Mandelstam variables s, $-t$ and $-u$ are large compared to a typical hadronic scale of order $\Lambda^2 = 1\,\mathrm{GeV}^2$. This implies $s, -t, -u \gg m^2$, where m is the proton mass. At Jefferson Lab (JLAB) there are ongoing experiments to measure the Compton cross section and certain spin transfer parameters [5]. Presently available beam energies are however not very high. In this note we investigate, as an example, the role of a non-negligible target mass in the handbag approach. An important issue in this context is the way to relate the dynamical variables of the approach to the external kinematics, determined by the experimental conditions. This relation is not unambiguous, which is one of the sources of theoretical uncertainties in the handbag approach. We study three different approximations and take the differences in their predictions as a measure of the theoretical uncertainty, which should be taken into account in attempts to extract the Compton form factors from experimental data.

The external kinematics is determined by the beam energy E_L^γ in the laboratory and by the scattering angle θ in the center of mass frame. These quantities fix the external Mandelstam variables by

$$\begin{aligned} s &= 2mE_L^\gamma + m^2\,, \\ t &= -\frac{s}{2}(1-\cos\theta)\,(1-m^2/s)^2\,, \\ u &= 2m^2 - s - t\,. \end{aligned} \tag{1}$$

These variables should not be changed or approximated in a theoretical calculation. Keeping this in mind we suggest a separate treatment of the kinematical factors from phase space and flux and of the scattering amplitude, which contains the dynamics. In the handbag approach the Compton cross section then reads [1, 3]

$$\begin{aligned} \frac{d\sigma}{dt} &= \frac{\pi\alpha_{\rm elm}^2}{(s-m^2)^2} \\ &\times \left[\frac{(\hat s-\hat u)^2}{|\hat s\hat u|}\Big(R_V^2(\hat t) - \frac{\hat t}{4m^2}R_T^2(\hat t)\Big) + \frac{(\hat s+\hat u)^2}{|\hat s\hat u|}R_A^2(\hat t)\right] \end{aligned} \tag{2}$$

to leading $\mathcal{O}(\alpha_s)$, where R_V, R_A and R_T are the Compton form factors. The one-loop corrections to the hard scattering have been evaluated in [3]. They were found to be small in the backward hemisphere and increased up to about 30% for $\cos\theta = 0.6$. The Mandelstam variables $\hat s$, $\hat t$, $\hat u$ refer to the partonic subprocess $\gamma q \to \gamma q$. They coincide with the external variables s, t, u up to corrections of order Λ^2/s. To calculate these consistently is beyond the accuracy of the approach in its present form. In particular, different choices for $\hat s$, $\hat t$, $\hat u$ lead to different results for the cross section at finite s.

We investigate the numerical effects of this ambiguity in three different scenarios. For the beam energy we take $E_L^\gamma = 4.3\,\mathrm{GeV}$, corresponding to $s = 8.97\,\mathrm{GeV}^2$, where there will soon be data from JLAB. We compare this with the situation for the energy $E_L^\gamma = 12\,\mathrm{GeV}$ of the proposed JLAB upgrade. We take the form factors R_V and R_A modeled in [1] using the overlap of light-cone wave functions and its connection to parton distributions and elastic form factors. From the overlap representation one expects a similar suppression of R_T/R_V as for the ratio F_2/F_1 of electromagnetic Pauli and Dirac form factors. For simplicity, we neglect R_T when evaluating the Compton cross section.

Scenario 1:

$$\hat s = s\,, \qquad \hat t = t\,, \qquad \hat u = u\,. \tag{3}$$

The subprocess amplitude leading to (3) was calculated in the approximation of massless on-shell

quarks, so that the sum of internal Mandelstam variables is zero. In this scenario this only holds approximately, since $\hat{s}+\hat{u}+\hat{t}=2m^2$.

Scenario 2:

$$\hat{s}=s-m^2\,,\qquad \hat{t}=t\,,\qquad \hat{u}=u-m^2\,, \tag{4}$$

where now $\hat{s}+\hat{u}+\hat{t}=0$.

Scenario 3:

$$\hat{s}=2m\,E_L^\gamma\,,\qquad \hat{t}=-\frac{\hat{s}}{2}\,(1-\cos\theta)\,,\qquad \hat{u}=-\hat{s}-\hat{t}\,. \tag{5}$$

Notice that in this case one has $\hat{t}\neq t$.

Numerical results for the Compton cross section evaluated from (3) without the R_T and α_s corrections are shown in Fig. 1(a) for the three scenarios. We plot the ratio

$$r=t^4\frac{d\sigma/dt}{d\sigma_{\rm KN}/dt}, \tag{6}$$

where

$$\frac{d\sigma_{\rm KN}}{dt} \;=\; \frac{2\pi\alpha_{\rm elm}^2}{(s-m^2)^2} \times\left[-\frac{s-m^2}{u-m^2}-\frac{u-m^2}{s-m^2}+\frac{4m^2t\,(m^4-su)}{(s-m^2)^2(u-m^2)^2}\right] \tag{7}$$

is the Klein-Nishina cross section for a point-like proton. The ratio r essentially measures an average of $(t^2R_V)^2$ and $(t^2R_A)^2$. These scaled form factors are expected to depend only weakly on t in the kinematical range considered here [1, 2].

For a beam energy of 4.3 GeV the differences between the cross sections evaluated in the three scenarios are moderate at small scattering angles but grows up to a factor 2 for backward angles. With a beam energy of 12 GeV instead, the ambiguities are small for all angles considered, as shown in Fig. 1(b).

In [1–3] we plotted the scaled Compton cross section $s^6d\sigma/dt$ with the squared proton mass neglected in both the internal and *external* variables, i.e. we used scenario 3 together with $s=\hat{s}$, $t=\hat{t}$, $u=\hat{u}$ and replaced $(s-m^2)^2$ with s^2 in (3). In this case the scattering angle θ and the scaling parameter s^6 multiplying the differential cross section do not correspond to the experimentally measured quantities. As a consequence both data and theoretical predictions in the plots of [1–3] were shifted. In other words we plotted $(2mE_L^\gamma)^6d\sigma/dt$ against $1+t/(mE_L^\gamma)$ rather than $s^6d\sigma/dt$ against $\cos\theta$. We realize that this is a rather confusing procedure which should be avoided in future presentations (we thank Travis Brooks and Lance Dixon for drawing our attention to this problem). Nevertheless, if our kinematical requirements of $s,-t,-u\gg\Lambda^2$ are well satisfied the expressions given in [1–3] are correct. The subtleties of internal vs. external Mandelstam variables matter only for energies as low as those currently available at JLAB, where the application of the handbag approach requires some care and is perhaps more qualitative.

Another interesting quantity is the correlation between the helicities of the incoming photon and the incoming (A_{LL}) or the outgoing (K_{LL}) proton in the c.m. In the handbag approach these parameters are given to leading $\mathcal{O}(\alpha_s)$ by [2, 3]

$$\frac{d\sigma}{dt}A_{LL}=\frac{d\sigma}{dt}K_{LL}=\frac{2\pi\alpha_{\rm elm}^2}{(s-m^2)^2} \times\;\frac{\hat{s}^2-\hat{u}^2}{|\hat{s}\hat{u}|}\,R_A(\hat{t})\left(R_V(\hat{t})+\frac{\hat{t}}{\hat{s}+\sqrt{-\hat{s}\hat{u}}}\,R_T(\hat{t})\right). \tag{8}$$

The corresponding expressions for a point-like proton are

$$\frac{d\sigma^{\rm KN}}{dt}\,A_{LL}^{\rm KN}=\frac{2\pi\alpha_{\rm em}^2}{(s-m^2)^2} \times\left[-\frac{s-m^2}{u-m^2}+\frac{u-m^2}{s-m^2}-\frac{2m^2t^2(s-u)}{(s-m^2)^2\,(u-m^2)^2}\right], \tag{9}$$

$$\frac{d\sigma^{\rm KN}}{dt}\,K_{LL}^{\rm KN}=\frac{2\pi\alpha_{\rm em}^2}{(s-m^2)^2} \times\left[-\frac{s-m^2}{u-m^2}+\frac{u-m^2}{s-m^2}-\frac{4m^2t^2(m^4-su)}{(s-m^2)^3\,(u-m^2)^2}\right],$$

which reduces to $A_{LL}^{\rm KN}=K_{LL}^{\rm KN}=(s^2-u^2)/(s^2+u^2)$ in the massless limit. In Fig. 2 we show the helicity correlation K_{LL} evaluated for the three scenarios, normalized to the Klein-Nishina value. This quantity is essentially a measure of R_A/R_V, as can be seen from

$$\frac{K_{LL}}{K_{LL}^{\rm KN}}\simeq\frac{R_A}{R_V}\,\left[1-\frac{t^2}{2(s^2+u^2)}\Big(1-\frac{R_A^2}{R_V^2}\Big)\right]^{-1}, \tag{10}$$

where we have neglected R_T and kinematic corrections of order m^2/s. Note that the kinematical prefactor in brackets is at most 0.3 for $\cos\theta\geq-0.6$ and $s\gg m^2$. Fig. 2 shows that for present JLAB energies the uncertainties related to the proton mass are sizeable, while at $E_L^\gamma=12$ GeV the effect is small.

JLAB will also measure the correlation K_{LS} between the helicity of the incoming photon and the sideways polarization of the outgoing proton. In the handbag approach it is given by [3]

$$\frac{d\sigma}{dt}K_{LS}=\frac{2\pi\alpha_{\rm elm}^2}{(s-m^2)^2} \times\;\frac{\hat{s}^2-\hat{u}^2}{|\hat{s}\hat{u}|}\,\frac{\sqrt{-\hat{t}}}{2m}\left(R_T(\hat{t})+\frac{4m^2}{\hat{s}+\sqrt{-\hat{s}\hat{u}}}\,R_V(\hat{t})\right)R_A(\hat{t}) \tag{11}$$

to leading $\mathcal{O}(\alpha_s)$, with the sign convention detailed in [3]. Contrary to the previous observables, K_{LS} is rather sensitive to the tensor Compton form factor R_T. It is convenient to consider the ratio K_{LS}/K_{LL} where in the

handbag approach the form factor R_A drops out. Introducing the abbreviation

$$\kappa_T(\hat{t}) \equiv \frac{\sqrt{-\hat{t}}}{2m}\,\frac{R_T(\hat{t})}{R_V(\hat{t})}\,, \tag{12}$$

this ratio can be expressed as [3]

$$\frac{K_{LS}}{K_{LL}} = \kappa_T\left(1+\frac{2m\sqrt{-\hat{t}}}{\hat{s}+\sqrt{-\hat{s}\hat{u}}}\,\frac{1}{\kappa_T}\right)\left(1-\frac{2m\sqrt{-\hat{t}}}{\hat{s}+\sqrt{-\hat{s}\hat{u}}}\,\kappa_T\right)^{-1}. \tag{13}$$

A rough estimate for the quantity κ_T can be obtained by considering the analogy between the ratio R_T/R_V and its electromagnetic counterpart F_2/F_1. One may, for instance, assume that

$$\kappa_T(\hat{t}) \simeq \frac{\sqrt{-\hat{t}}}{2m}\,\frac{F_2(\hat{t})}{F_1(\hat{t})} \approx 0.37\,, \tag{14}$$

where the numerical value on the r.h.s. is taken from the measurement of F_2/F_1 for $-t = 1 \div 5.6\,\mathrm{GeV}^2$ [6]. (Note, however, that on the basis of the previous SLAC measurement of F_2/F_1 [7] one would rather conclude that $\kappa_T \propto m/\sqrt{-t}$. For a detailed discussion of the uncertainties in the measurement of the Pauli form factor see [8].) We use (14) to estimate K_{LS} in the handbag approach and plot K_{LS}/K_{LL} for the three scenarios in Fig. 3. We observe that the ratio K_{LS}/K_{LL} is rather insensitive to the proton mass effects already at the present JLAB energy (in contrast, the predictions for K_{LS} alone suffer from the same uncertainties as for the other Compton observables discussed above). Measuring the ratio K_{LS}/K_{LL} at present JLAB energies, and solving for $\kappa_T(\hat{t})$ in (13) thus enables us to determine the ratio R_T/R_V and to test the analogy with F_2/F_1 in (14).

In conclusion, we found that finite proton mass effects severely limit the quantitative test of the handbag approach and the extraction of the Compton form factors in wide angle Compton scattering for present JLAB energies. An exception is the ratio K_{LS}/K_{LL} which turns out to be rather insensitive to finite proton mass effects, and can be used to determine the ratio of Compton form factors R_T/R_V. Qualitative features, like the sign and order of magnitude of the helicity correlations K_{LL} or K_{LS}, are not affected either. For higher photon energies as projected for JLAB the theoretical uncertainties from the proton mass become reasonably small.

[1] M. Diehl, T. Feldmann, R. Jakob and P. Kroll, Eur. Phys. J. C **8**, 409 (1999) [hep-ph/9811253].

[2] M. Diehl, T. Feldmann, R. Jakob and P. Kroll, Phys. Lett. B **460**, 204 (1999) [hep-ph/9903268].

[3] H. W. Huang, P. Kroll and T. Morii, Eur. Phys. J. C **23**, 301 (2002) [hep-ph/0110208].

[4] A. V. Radyushkin, Phys. Rev. D **58**, 114008 (1998) [hep-ph/9803316].

[5] J. P. Chen *et al.* [Jefferson Lab Hall A Collaboration], "Exclusive Compton Scattering On The Proton," Jefferson Lab preprint PCCF-RI-99-17.

[6] O. Gayou *et al.* [Jefferson Lab Hall A Collaboration], Phys. Rev. Lett. **88**, 092301 (2002) [nucl-ex/0111010].

[7] L. Andivahis *et al.*, Phys. Rev. D **50**, 5491 (1994).

[8] J. Arrington, hep-ph/0209243.



FIG. 1: (a) The ratio r defined in Eq. (6) at $E_L^\gamma = 4.3\,\mathrm{GeV}$ for scenarios 1 (full), 2 (long-dashed), 3 (short-dashed). The form factors R_V and R_A are taken from the model in [1] and R_T is neglected. (b) The same for $E_L^\gamma = 12\,\mathrm{GeV}$.



FIG. 2: (a) The ratio $K_{LL}/K_{LL}^{\mathrm{KN}}$ of helicity correlations at $E_L^\gamma = 4.3\,\mathrm{GeV}$ for scenarios 1 (full), 2 (long-dashed), 3 (short-dashed). R_V and R_A are taken from the model in [1] and R_T is neglected. (b) The same for $E_L^\gamma = 12\,\mathrm{GeV}$.



FIG. 3: (a) The ratio K_{LS}/K_{LL} of helicity correlations at $E_L^\gamma = 4.3\,\mathrm{GeV}$ for scenarios 1 (full), 2 (long-dashed), 3 (short-dashed). R_V is taken from the model in [1] and R_T is estimated from (14). (b) The same for $E_L^\gamma = 12\,\mathrm{GeV}$.